



AAPC-CPC^{Q&As}

Certified Professional Coder (CPC) Exam

Pass AAPC AAPC-CPC Exam with 100% Guarantee

Free Download Real Questions & Answers **PDF** and **VCE** file from:

<https://www.geekcert.com/aapc-cpc.html>

100% Passing Guarantee
100% Money Back Assurance

Following Questions and Answers are all new published by AAPC
Official Exam Center

-  **Instant Download** After Purchase
-  **100% Money Back** Guarantee
-  **365 Days** Free Update
-  **800,000+** Satisfied Customers





QUESTION 1

View MR 002395 MR 002395 Operative Report Pre-operative Diagnosis: Acute rotator cuff tear Post-operative Diagnosis: Acute rotator cuff tear, synovitis Procedures: 1) Rotator cuff repair 2) Biceps Tenodesis 3) Claviclectomy 4) Coracoacromial ligament release Indication: Rotator cuff injury of a 32-year-old male, sustained while playing soccer. Findings: Complete tear of the right rotator cuff, synovitis, impingement. Procedure: The patient was prepared for surgery and placed in left lateral decubitus position. Standard posterior arthroscopy portals were made followed by an anterior- superior portal. Diagnostic arthroscopy was performed. Significant

synovitis was carefully debrided. There was a full-thickness upper 3rd subscapularis tear, which was repaired. The lesser tuberosity was debrided back to bleeding healthy bone and a Mitek 4.5 mm helix anchor was placed in the lesser tuberosity. Sutures were passed through the subcapulans in a combination of horizontal mattress and simple interrupted fashion and then tied. There was a partial-thickness tearing of the long head of the biceps. The biceps were released and then anchored in the intertubercular groove with a screw. There was a large anterior acromial spur with subacromial impingement. A CA ligament was released and acromioplasty was performed. Attention was then directed to the

supraspinatus tendon tear. The tear was V-shaped and measured approximately 2.5 cm from anterior to posterior. Two Smith and Nephew PEEK anchors were used for the medial row utilizing Healicoil anchors. Side-to-side stitches were placed. One set of suture tape from each of the medial anchors was then placed through a laterally placed Mitek helix PEEK knotless anchor which was fully inserted after tensioning the tapes. A solid repair was obtained. Next there were severe degenerative changes at the AC joint of approximately 8 to 10 mm. The distal clavicle was resected taking care to preserve the superior AC joint capsule. The shoulder was thoroughly lavaged. The instruments were removed and the incisions were closed in routine fashion. Sterile dressing was applied. The patient was transferred to recovery in stable condition.

What CPT coding is reported for this case?

- A. 29827, 29828-51, 29824-51, 29826
- B. 29827, 29824-51, 29826-51
- C. 29827, 29828-51, 29824-51, 29826, 29805-59
- D. 29827, 29824-51, 29826-51, 29805-59

Correct Answer: A

QUESTION 2

The outermost protective layer of skin is called the:

- A. Epidermis
- B. Hypodermis
- C. Subcutaneous tissue
- D. Dermis

Correct Answer: A



QUESTION 3

A 47-year-old female presents to the operating room for a partial corpectomy on one upper thoracic vertebral body, T3. Two surgeons are performing the surgery. One surgeon performs the transthoracic approach and excises the damaged portion of the vertebral body. The second surgeon inserts a bone graft into the vertebral gap, closing the gap, and inserts a metal plate. Both surgeons work together, each as a primary surgeon.

How does each surgeon report their portion of the surgery?

- A. 63090-66, 63091-66
- B. 63087-62, 63088-62
- C. 63090-80, 63091-80
- D. 63085-62, 63086-62

Correct Answer: D

QUESTION 4

Which entity offers compliance program guidance to form the basis of a voluntary compliance program for a provider practice?

- A. Centers for Medicare and Medicaid Services (CMS)
- B. American Medical Association (AMA)
- C. Office of Inspector General (OIG)
- D. Office for Civil Rights (OCR)

Correct Answer: C

QUESTION 5

A 44-year-old female patient with chest pains had a CT of her chest that identified a mass in her left lower lung. The patient currently has ovarian cancer with metastases to the liver. The radiologist suspects the cancer has spread to her lungs. The physician performed an outpatient bronchoscopic biopsy and the pathology report documents the mass as a tumor of uncertain behavior.

What ICD-10-CM codes are reported for this patient?

- A. R91.8, C56.9, C78.7
- B. C56.9, C78.7, C78.02
- C. C78.02, C22.9, C79.82
- D. D38.1, C56.9, C78.7

Correct Answer: A



VCE & PDF

GeekCert.com

<https://www.geekcert.com/aapc-cpc.html>

2024 Latest geekcert AAPC-CPC PDF and VCE dumps Download

[Latest AAPC-CPC Dumps](#)

[AAPC-CPC VCE Dumps](#)

[AAPC-CPC Practice Test](#)