



NAPLEX^{Q&As}

North American Pharmacist Licensure Examination

Pass Test Prep NAPLEX Exam with 100% Guarantee

Free Download Real Questions & Answers **PDF** and **VCE** file from:

<https://www.geekcert.com/naplex.html>

100% Passing Guarantee
100% Money Back Assurance

Following Questions and Answers are all new published by Test Prep
Official Exam Center

-  **Instant Download** After Purchase
-  **100% Money Back** Guarantee
-  **365 Days** Free Update
-  **800,000+** Satisfied Customers





QUESTION 1

If a patient is getting NS IVF at 120mls/hr, how much Sodium Chloride in grams is this patient getting in 24 hours? NS is 0.9% sodium chloride.

- A. 25.92gm
- B. 25.92kg
- C. 25.92mg
- D. 90mg
- E. 90gm

Correct Answer: A

QUESTION 2

A 54-year-old male with a long history of mild persistent asthma on daily fluticasone therapy has been using his albuterol inhaler every day for the past month, and presents requesting a refill. What changes should be made to his current regimen?

- A. Add ciclesonide to current regimen
- B. Add salmeterol to current regimen
- C. Discontinue fluticasone and instead use salmeterol
- D. Add cromolyn to current regimen
- E. Discontinue fluticasone and add ipratropium to current regimen

Correct Answer: B

Add salmeterol to the current regimen. This patient had mild persistent asthma but was using his albuterol daily, which indicates that a step up in therapy is warranted. The preferred first line treatment regimen for moderate persistent asthma are low to medium dose inhaled corticosteroids plus a long acting beta2 agonist, as well as a short acting beta2 agonist as needed. A is incorrect. Ciclesonide is an inhaled corticosteroid. The patient in the case is already using fluticasone, so adding ciclesonide would be therapeutic duplication. C is incorrect. Long-acting beta2 agonists should only be used as adjunctive therapy in patients who are currently receiving but not adequately controlled on an inhaled corticosteroid. These medications should not be used as monotherapy, due to an increased risk of asthma related deaths. D is incorrect. Cromolyn prevents the release of vasoactive mediators from mast cell and is primarily used for exercise-induced asthma, it is not indicated as an alternative agent in patients with moderate persistent asthma. E is incorrect. Ipratropium is a short-acting anticholinergic, which is often used in COPD or in asthma exacerbations. It is not indicated for maintenance treatment of moderate persistent asthma.

QUESTION 3



According to the 2013 ACC/AHA Guideline on the Treatment of Blood Cholesterol to Reduce Atherosclerotic Cardiovascular Risk in Adults, which of the following medications and dosing regimens is considered moderate-intensity statin therapy?

- A. Atorvastatin 80 mg PO QHS
- B. Lovastatin 20 mg PO QHS
- C. Simvastatin 40 mg PO QHS
- D. Rosuvastatin 40 mg PO QHS
- E. Atorvastatin 20mg PO QHS

Correct Answer: E

QUESTION 4

Before use, which of the following products should be used to clean surfaces of a laminar flow hood?

- A. 95% ethyl alcohol
- B. 70% isopropyl alcohol
- C. Purified water
- D. Purified WFI

Correct Answer: B

Before each use, 70% isopropyl alcohol should be used to clean surfaces of a laminar flow hood.

QUESTION 5

What is the standard loading dose for clopidogrel in the treatment of ACS?

- A. 75mg
- B. 100mg
- C. 150mg
- D. 200mg
- E. 300mg

Correct Answer: E

300mg is the standard loading dose of clopidogrel ?an antiplatelet drug ?in the treatment of acute coronary syndrome (ACS).



VCE & PDF

GeekCert.com

<https://www.geekcert.com/naplex.html>

2024 Latest geekcert NAPLEX PDF and VCE dumps Download

[Latest NAPLEX Dumps](#)

[NAPLEX PDF Dumps](#)

[NAPLEX Practice Test](#)